



AIP Scilight

HOME

BROWSE

INFO



12 FEBRUARY 2018 •

<https://doi.org/10.1063/1.5025497>

1952 physics model describes feather bud formation observed in vitro


Randolph Fillmore

Researchers investigate role of boundaries and self-organizing processes in feather bud formation.

206
VIEWS

12
SHARES





↓ PPT | High-resolution

Skin self-organization and natural, “genetically mapped” boundaries may be an important feature in bioengineering and regenerating skin appendages, such as feathers. Like other skin appendages, teeth, hair or even lung tissue, feathers are developed via interactions between the epithelium and the mesenchyme – mesodermal embryonic tissue – during early stages of morphogenesis. Authors of an article in *APL Bioengineering* describe their experiment evaluating pattern formation in feather buds growing in vitro, employing Turing’s model, a “two reaction-diffusion model” published in 1952. The work advances the

hypothesis that patterns during embryonic development respond to spatial biochemical pre-patterning.

Turing hypothesized that patterns seen in nature reflect underlying biochemical signaling. Similarly, the researchers found that a boundary effect guided the formation and patterning of feather buds in chicks. The formation they used its numerical description used a standard two-component reaction model.

Feather bud morphological features changed dramatically during the first two days of incubation as well as from days 0-4. Prime locations for feather bud formation were near the skin edges, a boundary edge that may function as a wall guiding the self-organization as bud formations occur at predictable sites from skin edges.

Co-author Toshiyuki Mitsui said their results revealed high sensitivity to the size and shape of edges of bioengineered tissue composed of uniformly distributed homogeneous cells. This observation, he said, provides potential templates for studying the self-organizing processes in advancing regenerative medicine.

In the future, they will try to grow human hair using induced pluripotent stem cells (iPSCs), stem cells that can be generated directly from adult cells. The researchers said there may be an optimized geometry for each organ to regenerate from iPSCs.

Source: “Role of the boundary in feather bud formation on one-dimensional bioengineered skin,” by Kentaro Ishida and Toshiyuki Mitsui,

APL Bioengineering (2018). The article can be accessed at

<https://doi.org/10.1063/1.4989414>.

Published by AIP Publishing (<https://publishing.aip.org/authors/rights-and-permissions>).

MOST READ

JUL 24 2017

Magnetic switching of ferromagnetic materials with a single femtosecond laser pulse

JUN 27 2017

New study provides guide on how to make your own hyperuniform system

AUG 21 2017

Novel poling method in piezoelectric ceramics opens the door to the next-generation large-strain actuators

SEP 01 2017

Extending the large bubble entrapment regime and modelling the mechanics

AUG 25 2017

Using the Cohen-Procaccia function to extract entropy origin in photonic random number generators

Subscribe to Scilight Alerts!

Receive weekly updates with the most interesting articles

SUBSCRIBE

Resources

AUTHOR

LIBRARIAN

ADVERTISER

General Information

ABOUT

[CONTACT](#)

[HELP](#)

[PRIVACY POLICY](#)

[TERMS OF USE](#)

FOLLOW AIP PUBLISHING:



Website © 2018 AIP Publishing LLC. Article copyright remains as specified within the article.

Scitation

[← PREV](#)